	Sector: Poultry					
	Date: 28/07/2020					
	☑ Meeting ☐ tel ☐ e-mail					
	Participants: Olivier SALANDRE (SNGTV Veterinary Representative) Claire CHAUVIN (Anses Ploufragan) - in Skype					
	New/ added comments post meeting					
	General remarks: The deficiency in terms of antipyretics, analges The New Regulation will change the fixed withdrawal times appl (reminder: the oldest marketing authorisations are suitable for withdrawal time).	licable under the "cascade" in a number of cases: commun	ication on change	es with a significa		mpact would be helpful. C leads to difficulties of use under the "cascade" due to the imposed
Meeting PRIORITIES			l			
Major: M minor: m	Risks	Action to be taken	Who	Deadline	Status	Comments
	OTC use on viral diseases. Risk of using aspirin or paracetamol in absence of MRL dossier on eggs and poultry. Acetyl salicylic acid (="aspirin"), sodium acetylsalicylate and sodium salicylate: "Do not use in animals producing milk or eggs intended for human consumption". For sodium salicylate, the EPMAR of 15 January 2013 (EMA/CVMP/71291/ 2012) on the extension of MRLs to turkeys specifies that the active substance must not be used in animals producing eggs intended for human consumption. The EPMAR of 12 December 2016 (EMA/CVMP/454104/2016) specifies that there is no extrapolation possible from turkey MRLs to broilers and laying hens. The NVR will confirm the ban on poultry use via the cascade. Nothing in pain management in laying machines (Animal welfare problem).	body. (For sodium salicylate, with the MRLs defined for turkey	Industrial via SIMV Anses Ploufragan	To be specified		According to regulation 2017/880, no extrapolation possible between unrelated species, or between tissues and eggs within the same species Need for data on metabolism and residues in poultry to obtain an MRL for eggs. It should be noted that the NVR will support this prohibition of use in the absence of data and ensure 5 years protection on the data submitted.
M (but no short- term solution)		Possibility of setting up studies with possible funding from the sector in conjunction with what is done in Austria.	Anses Ploufragan	To be completed	In progress	
M (multi-channel transversal priority)	Animal welfare problem and societal expectations	* Discuss identified needs with Anses DER	Anses Ploufragan ANMV			
	-					

m	Use of Colistin and Aminoglycosides against E.coli, while not	* Prevention method: background study to be conducted on the epidemiology of colibacilli, vertical/horizontal	Sector &	2022		
	absorbed by the digestive tract => inappropriate use ("disinfectant" type).	contamination.	Anses Ploufragan			
	(districtant type).	Contamination	ricanagan			
		* Biosecurity: propose actions as part of the Eco antibio	Sector?			
		2 programme (DGAL funding)?				
		* Inform about CVMAD modifications on Envallages in	ANIBA)/			Indication withdrawn since oninion of the CVAAD of Feb. 2019
		* Inform about CVMP modifications on Enrofloxacin indications	ANMV			Indication withdrawn since opinion of the CVMP of Feb. 2018
		* Inform about the ability to produce autovaccines	MAH		Completed	Advertising is prohibited for auto-vaccines in the decree and in the future
		without mentioning the target strains/species				regulations.
		(depending on interest)				However, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they
						are authorised.
						are dutionsed.
		* Poulvac E.coli: provide studies for laying extension to laying hens (currently "contraindications")?	MAH (Zoétis)	To be specified		
		laying nens (currently contramucations):				
	Non-compliance of WT when using toltrazuril in standard	To be specified				
m	broilers.	To be specified				
	"Phyto" alternatives used without an official claim and rather					
	indications such as "contributes to the management of the risk					
	associated with coccidia". Products used in drinking water, therefore equivalent to treatments. No rigorous efficacy					
	studies.					
	harmonist and off label are of flaboured	* Boundary and the United States and another the Control of the Co	to disease to 1	To be seed 0.1		Description and trade and to Table 4 with an AADI appropriate form.
m	Inappropriate and off-label use of flubendazole or fenbendazole at higher or more repeated doses	* Develop metabolism data and residues in poultry to obtain MRL Praziquantel poultry eggs (and meat).	Industrial via SIMV	To be specified		Praziquantel included in Table 1 with no MRL required for sheep (total residues < 30% ADI) and horse.
	Temberidazore at filigiler of more repeated doses	* Plan field trials (on breeders for example) to study the	VIA SIIVIV			residues \ 30% ADIJ and norse.
	=> impact on residues/environment?	efficacy of praziquantel.	Sector			=> Need for metabolism data and poultry residues to be provided for possible
						extrapolation to this species.
		* What is the prevalence of teniasis in outdoor farming?	Anses		Completed	Helminth Observatory set up in 2019-20 by ITAVI & Anses (75% funded by
			Ploufragan			pharmaceutical laboratories and 25% by production organisations)
			Sector			=> Information on diagnostic methods: see article published in the journal
			Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-des-
			Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-des- tamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles)
			Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-des-
			Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-yers-parasites-des-volailles) => A publication in TeMa is planned, either end of year or early next year (1st
			Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-yers-parasites-des-volailles) => A publication in TeMa is planned, either end of year or early next year (1st
			Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-yers-parasites-des-volailles) => A publication in TeMa is planned, either end of year or early next year (1st
m	Currently MRL enilconazole only for topical use while the	* Rework the IMAVERAL file for turkeys to avoid the	Sector			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-yers-parasites-des-volailles) => A publication in TeMa is planned, either end of year or early next year (1st
m	Currently MRL enilconazole only for topical use while the product is used against aspergillosis by spraying in the	* Rework the IMAVERAL file for turkeys to avoid the cascade.				⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results.
m			МАН			=> Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-evers-parasites-des-volailles) => A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys.
m 0	product is used against aspergillosis by spraying in the	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be	МАН		Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in
	product is used against aspergillosis by spraying in the	cascade. *What are the risks for the user of off-label spray use?	MAH (Elanco)		Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-det-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested?
0	product is used against aspergillosis by spraying in the	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)?	MAH (Elanco) ANMV	Tobocasio	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-det-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl)
	product is used against aspergillosis by spraying in the	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? * Inform about the ability to produce autovaccines	MAH (Elanco)	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future
0	product is used against aspergillosis by spraying in the	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)?	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-det-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl)
0	product is used against aspergillosis by spraying in the	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? * Inform about the ability to produce autovaccines without mentioning the target strains/species	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for
0	product is used against aspergillosis <u>by spraying in the</u> <u>presence of animals.</u>	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? * Inform about the ability to produce autovaccines without mentioning the target strains/species	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-det-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to
0	product is used against aspergillosis by spraying in the	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for
0	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What are the risks for the user of off-label spray use? *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? * Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest)	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for
0	product is used against aspergillosis <u>by spraying in the</u> <u>presence of animals.</u>	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for
0	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for
0	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in	MAH (Elanco) ANMV	To be specified	Completed	⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for
m m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in turkeys?	MAH (Elanco) ANMV Owners String?	To be specified		⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised.
m m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in turkeys?	MAH (Elanco) ANMV Owners String?	To be specified		⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-de-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain
m m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in turkeys?	MAH (Elanco) ANMV Owners String?	To be specified 31/07/20		⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-de-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain
m m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? * Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) * Do we have information on the use of dimetridazole in turkeys? *What is the impact of NVR on WT to be applied?	MAH (Elanco) ANMV Owners String?			⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-de-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain antibiotics Check if marketing of the smallest packs available: COLOMBOVAC PMV (Zoétis) - pigeons: 1 vial of 10 doses, 50 doses -
0 m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) *Do we have information on the use of dimetridazole in turkeys? *What is the impact of NVR on WT to be applied? *Information on availability of small packs of vaccines	MAH (Elanco) ANMV Owners String?			⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-de-tquantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain antibiotics Check if marketing of the smallest packs available: COLOMBOVAC PMV (Zoétis) - pigeons: 1 vial of 10 doses, 50 doses - unavailable?
0 m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) *Do we have information on the use of dimetridazole in turkeys? *What is the impact of NVR on WT to be applied? *Information on availability of small packs of vaccines	MAH (Elanco) ANMV Owners String?			⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain antibiotics Check if marketing of the smallest packs available: COLOMBOVAC PMV (Zoétis) - pigeons: 1 vial of 10 doses, 50 doses - unavailable? COBILIS PARAMYXO P201 (Intervet) - pigeons: 1 vial of 20 mL (=80 doses)
m m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) *Do we have information on the use of dimetridazole in turkeys? *What is the impact of NVR on WT to be applied? *Information on availability of small packs of vaccines	MAH (Elanco) ANMV Owners String?			⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-de-t-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fowl) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain antibiotics Check if marketing of the smallest packs available: COLOMBOVAC PMV (Zoétis) - pigeons: 1 vial of 10 doses, 50 doses - unavailable? NOBILIS PARAMYYO P201 (Intervet) - pigeons: 1 vial of 20 mL (=80 doses) IMOPEST (BI) - chickens & pigeons: 1 vial of 500 doses
0 m	product is used against aspergillosis <u>by spraying in the presence of animals.</u> Risk of using prohibited products (dimetridazole). ???	cascade. *What is the impact of NVR on WT parconazole to be applied to turkey (0 days in guinea fowl)? *Inform about the ability to produce autovaccines without mentioning the target strains/species (depending on interest) *Do we have information on the use of dimetridazole in turkeys? *What is the impact of NVR on WT to be applied? *Information on availability of small packs of vaccines	MAH (Elanco) ANMV Owners String?			⇒ Information on diagnostic methods: see article published in the journal TeMA and video tutorials (e.g. https://www.reussir.fr/volailles/utiliser-destamis-pour-identifier-et-quantifier-les-vers-parasites-des-volailles) ⇒ A publication in TeMa is planned, either end of year or early next year (1st quarter) which will present the details of the results. Nothing new. MRL file to be developed for Turkeys. MRL exists for cattle (total residues < 11% ADI) and horses for topical use. To be requested? Parconazole BP to be applied to turkey: with NVR, change to 1 day (0 days in guinea fow!) Advertising is prohibited for autovaccines in the decree and in the future regulations. On the other hand, it may be possible to communicate about the ability to produce autovaccines without mentioning the target strains/species for which they are authorised. NVR impact on cascade may allow more favourable WTs for certain antibiotics Check if marketing of the smallest packs available: COLOMBOVAC PMV (Zoétis) - pigeons: 1 vial of 10 doses, 50 doses - unavailable? COBILIS PARAMYXO P201 (Intervet) - pigeons: 1 vial of 20 mL (=80 doses)